

Energy distribution of the Einstein-Klein-Gordon system for a static spherically symmetric spacetime in $(2 + 1)$ -dimensions

I. Radinschi^{*1}, Th. Grammenos^{†2}

** Physics Department, “Gh. Asachi” Technical University,
6600 Iasi, Romania*

*† Department of Mechanical & Industrial Engineering, University of Thessaly,
383 34 Volos, Greece*

Abstract

We use Møller’s energy-momentum complex in order to explicitly compute the energy and momentum density distributions for an exact solution of Einstein’s field equations with a negative cosmological constant minimally coupled to a static massless scalar field in a static, spherically symmetric background in $(2 + 1)$ -dimensions.

Keywords: Energy-momentum complex, scalar field

PACS numbers: 04.20.-q, 04.20.Jb

¹radinschi@yahoo.com

²thgramme@uth.gr

1 Introduction

One of the most interesting issues in General Relativity is the subject of energy-momentum localization. This problem of defining in an acceptable manner the energy-momentum density hasn't got a generally accepted answer yet. The localization of energy implies the use of various energy-momentum complexes, including the prescriptions of Einstein [1], Landau and Lifshitz [2], Papapetrou [3], Bergmann [4] and Weinberg [5]. All of these prescriptions have a drawback: the calculations are restricted to quasi-Cartesian coordinates. On the other hand, Møller [6] proposed an energy-momentum complex which enables one to calculate the energy distribution in any coordinate system and not only in quasi-Cartesian coordinates. Hence, an interesting question is whether any of the aforementioned prescriptions provides the best option for energy-momentum localization and, of course, under what conditions.

The problem of energy-momentum localization by using the energy-momentum complexes was revived by Virbhadra and his collaborators. He established an important result showing that different energy-complexes (ELLPW) yield the same result for a general non-static spherically symmetric metric of the Kerr-Schild class [7]. Furthermore, these definitions (ELLPW) comply with the quasi-local mass definition of Penrose for a general non-static spherically symmetric metric of the Kerr-Schild class. However, these prescriptions disagree in the case of the most general non-static spherically symmetric metric. According to Virbhadra, although the energy-momentum complexes behave under general coordinate transformations like non-tensorial objects, the local conservation laws obtained by them hold in all coordinate systems. Also, many interesting results recently obtained [8]-[9] point out that the energy-momentum complexes are powerful tools for obtaining the energy distribution in a given spacetime. Interesting works were done with the energy-momentum complexes in 2- and 3-dimensional spacetimes [10]. In this context, of great importance is the Cooperstock hypothesis [11] which states that energy and momentum are confined to the regions of non-vanishing energy-momentum tensor for the matter and all non-gravitational fields.

Chang, Nester and Chen [12] showed that the energy-momentum complexes are actually quasilocal and give a legitimate expression for the energy-momentum. Further, they concluded that there exists a direct relationship between energy-momentum complexes and quasilocal expressions, since every energy-momentum complex is associated with a legitimate Hamiltonian boundary term.

Regarding Møller's energy-momentum complex, there is a plethora of results [13]-[14] that recommend it as an efficient tool for the localization of energy. According to Lessner [15], Møller's energy-momentum complex is significant for describing the concepts of energy and momentum in General Relativity. In fact, he stated that "The energy-momentum four-vector can transform according to special relativity only if it is transformed to a reference system with an everywhere constant velocity. This cannot be achieved by a global Lorentz transformation".

For Bondi [16] "a nonlocalizable form of energy is not admissible in General Relativity, because any form of energy contributes to gravitation and so its location can in principle

be found”.

Based on the above works and conclusions we have decided to use Møller’s energy-momentum complex for evaluating the energy distribution of an exact solution for a static massless scalar field minimally coupled to gravity in $(2 + 1)$ dimensions.

Consequently, in this work we have implemented Møller’s formalism to calculate the energy distribution for a geometry given recently by Daghan and Bilge [17]. They used a series of transformations to reduce to second order the system of differential equations for the Einstein’s field equations with a negative cosmological constant, minimally coupled to a massless scalar field in a static, spherically symmetric background in $(2+1)$ -dimensions. In this way, the authors obtained an exact solution in a form that generalizes the AdS metric as given by Matschull [18]. Throughout we use geometrized units, i.e. $c = G = 1$, while $g_{\mu\nu}$ has the signature $(-, +, +)$, Greek indices run from 0 to 2 and Latin indices run over the spatial coordinate values. The rest of the paper is organized as follows. In Sec.2 we give a short review of the concept of energy-momentum complexes and present Møller’s prescription. In Sec.3 we briefly present the $(2 + 1)$ -dimensional, spherically symmetric background used, while in Sec.4 we explicitly compute the energy and momentum density distributions of the Einstein-Klein-Gordon system using Møller’s energy-momentum complex. Finally, in Sec.5 a discussion of the obtained results and some concluding remarks are given.

2 Energy-Momentum Complexes and Møller’s Prescription

The conservation laws of energy and momentum for an isolated, i.e., no external force is acting on the system, physical system in the Theory of Special Relativity are expressed by a set of differential equations. Defining T^μ_ν as the symmetric energy-momentum tensor of matter and of all non-gravitational fields, the conservation laws are given by

$$T^\mu_{\nu,\mu} \equiv \frac{\partial T^\mu_\nu}{\partial x^\mu} = 0, \quad (2.1)$$

where

$$\rho = T^t_t \quad j^i = T^i_t \quad p_i = -T^t_i \quad (2.2)$$

are the energy density, the energy current density, and the momentum density, respectively. Making the transition from Special to General Theory of Relativity, one adopts a simplicity principle which is called principle of minimal gravitational coupling. As a result, the conservation equation is now written as

$$T^\mu_{\nu;\mu} \equiv \frac{1}{\sqrt{-g}} \frac{\partial}{\partial x^\mu} (\sqrt{-g} T^\mu_\nu) - \Gamma^\kappa_{\nu\lambda} T^\lambda_\kappa = 0, \quad (2.3)$$

where g is the determinant of the metric tensor $g_{\mu\nu}(x)$. The conservation equation may also be written as

$$\frac{\partial}{\partial x^\mu} (\sqrt{-g} T^\mu_\nu) = \xi_\nu, \quad (2.4)$$

where

$$\xi_\nu = \sqrt{-g} \Gamma_{\nu\lambda}^\kappa T_\kappa^\lambda \quad (2.5)$$

is a non-tensorial object. For $\nu = t$ this means that the matter energy is not a conserved quantity for the physical system³. From a physical point of view, this lack of energy conservation can be understood as the possibility of transforming matter energy into gravitational energy and vice versa. However, this remains an open problem and it is widely believed that in order to solve it one has to take into account the gravitational energy.

By a well-known procedure, the non-tensorial object ξ_ν can be written as

$$\xi_\nu = -\frac{\partial}{\partial x^\mu} (\sqrt{-g} \vartheta_\nu^\mu), \quad (2.6)$$

where ϑ_ν^μ are certain functions of the metric tensor and its first order derivatives. Therefore, the energy-momentum tensor of matter T_ν^μ is replaced by the expression

$$\theta_\nu^\mu = \sqrt{-g} (T_\nu^\mu + \vartheta_\nu^\mu), \quad (2.7)$$

which is called energy-momentum complex since it is a combination of the tensor T_ν^μ and a pseudotensor ϑ_ν^μ describing the energy and momentum of the gravitational field. The energy-momentum complex satisfies a conservation law in the ordinary sense, i.e.,

$$\theta_{\nu,\mu}^\mu = 0 \quad (2.8)$$

and it can be written as

$$\theta_\nu^\mu = \chi_{\nu,\lambda}^{\mu\lambda}, \quad (2.9)$$

where $\chi_{\nu,\lambda}^{\mu\lambda}$ are called superpotentials and are functions of the metric tensor and its first order derivatives.

It is evident that the energy-momentum complex is not uniquely determined by the condition that its usual divergence is zero since a quantity with an identically vanishing divergence can always be added to the energy-momentum complex.

The energy-momentum complex of Møller in a four-dimensional background is given as [6]

$$\mathcal{J}_\nu^\mu = \frac{1}{8\pi} \xi_{\nu,\lambda}^{\mu\lambda}, \quad (2.10)$$

where Møller's superpotential $\xi_{\nu,\lambda}^{\mu\lambda}$ is of the form

$$\xi_{\nu,\lambda}^{\mu\lambda} = \sqrt{-g} \left(\frac{\partial g_{\nu\sigma}}{\partial x^\kappa} - \frac{\partial g_{\nu\kappa}}{\partial x^\sigma} \right) g^{\mu\kappa} g^{\lambda\sigma}, \quad (2.11)$$

with the antisymmetric property

$$\xi_{\nu,\lambda}^{\mu\lambda} = -\xi_{\nu,\lambda}^{\lambda\mu}. \quad (2.12)$$

³It is possible to restore the conservation law by introducing a local inertial system for which at a specific spacetime point $\xi_\nu = 0$ but this equality by no means holds in general.

It is easily seen that Møller's energy-momentum complex satisfies the local conservation equation

$$\frac{\partial \mathcal{J}_\nu^\mu}{\partial x^\mu} = 0, \quad (2.13)$$

where $\mathcal{J}_0^0$ is the energy density and $\mathcal{J}_i^0$ are the momentum density components.

Thus, in Møller's prescription the energy and momentum for a four-dimensional background are given by

$$P_\mu = \int \int \int \mathcal{J}_\mu^0 dx^1 dx^2 dx^3. \quad (2.14)$$

Specifically the energy of the physical system in a four-dimensional background is

$$E = \int \int \int \mathcal{J}_0^0 dx^1 dx^2 dx^3. \quad (2.15)$$

It should be noted that the calculations are not anymore restricted to quasi-Cartesian coordinates but can be utilized in any coordinate system.

3 The (2 + 1)-dimensional Static Spherically Symmetric Background

The line element of a three-dimensional static, spherically symmetric, Lorentzian space-time is given as

$$ds^2 = -e^{2f(r)} dt^2 + e^{2g(r)} dr^2 + e^{-2h(r)} d\theta^2, \quad (3.1)$$

in (t, r, θ) coordinates. Einstein's field equations with a cosmological constant Λ are

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu}, \quad (3.2)$$

where

$$T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} g_{\mu\nu} (\partial_\lambda \phi \partial^\lambda \phi) \quad (3.3)$$

is the energy-momentum tensor of a massless scalar field Φ [19] satisfying the wave equation:

$$g^{\mu\nu} \nabla_\mu \nabla_\nu \phi = (-g)^{-\frac{1}{2}} \partial_\mu [(-g)^{-\frac{1}{2}} g^{\mu\nu} \partial_\nu \phi]. \quad (3.4)$$

The above wave equation of the Klein-Gordon type can be integrated giving for the first derivative of the static scalar field ϕ :

$$\phi' = \frac{\lambda}{\sqrt{2\pi}} e^{h(r)}, \quad (3.5)$$

where λ is an integration constant. By using a series of transformations and taking $\Lambda = -1$, Daghan and Bilge [17] managed to solve analytically the above Einstein-Klein-Gordon system of differential equations for $f(r)$, $h(r)$ and ϕ and obtained the following form for the line element:

$$ds^2 = - \left[\frac{\cosh \chi}{\sinh \chi} \right]^\mu \cosh \chi \sinh \chi d\tilde{t}^2 + d\chi^2 + (c^2 + 4\lambda^2) \left[\frac{\cosh \chi}{\sinh \chi} \right]^{-\mu} \cosh \chi \sinh \chi d\tilde{\theta}^2, \quad (3.6)$$

where $\chi, \tilde{t}, \tilde{\theta}$ are defined through the following transformations:

$$\chi = -\frac{1}{2} \ln(\tan \frac{\varphi}{2}), \quad (3.7)$$

$$\tilde{t} = \frac{\rho_0}{2} t, \quad (3.8)$$

$$\tilde{\theta} = \frac{2}{\rho_0} \theta. \quad (3.9)$$

$c \neq 0$ being an integration constant and $\mu = \frac{c}{\sqrt{c^2 + 4\lambda^2}}$.

$\varphi \in (0, \pi/2)$ and it can be seen that for $\varphi = \pi/2$ a curvature singularity arises. Furthermore, φ is related to $f(r)$ and $h(r)$ through

$$e^{f(r)} = \frac{1}{2} \rho \cos \varphi, \quad (3.10)$$

$$e^{h(r)} = \frac{1}{\sqrt{c^2 + 2\lambda^2}} \rho \sin \varphi, \quad (3.11)$$

while

$$\rho(\varphi) = \frac{\rho_0}{\sqrt{2}} (1 + \sin \varphi)^{\frac{\mu}{2}} (\sin \varphi)^{-\frac{1}{2}} (\cos \varphi)^{\frac{-1-\mu}{2}} \quad (3.12)$$

and ρ_0 is an integration constant too.

It should be noted, that for $\lambda = 0$ and $c = 1$ one gets the three-dimensional Anti-deSitter solution given by Matschull in [18].

4 Energy and Momentum Density Distributions

We first have to evaluate the superpotentials in the context of Møller's prescription for the spacetime described by the line element (3.6). There are four non-zero independent superpotentials:

$$\begin{aligned} \xi_0^{01} &= (-\mu + \cosh(2\chi)) \sqrt{c^2 + 4\lambda^2}, \\ \xi_0^{10} &= -\xi_0^{01} = (\mu - \cosh(2\chi)) \sqrt{c^2 + 4\lambda^2}, \\ \xi_2^{21} &= \frac{(9 + 4\lambda^2)(\mu + \cosh(2\chi))}{\sqrt{c^2 + 4\lambda^2}}, \\ \xi_2^{12} &= -\xi_2^{21} = -\frac{(9 + 4\lambda^2)(\mu + \cosh(2\chi))}{\sqrt{c^2 + 4\lambda^2}}. \end{aligned} \quad (4.1)$$

Substitution of the above superpotentials into eq.(2.10) yields the energy density distribution

$$\mathcal{J}_0^0 = \frac{4}{K} \sqrt{c^2 + 4\lambda^2} \cosh \chi \sinh \chi, \quad (4.2)$$

where K is an integration constant, while the momentum density distributions obtained are

$$\mathcal{J}_1^0 = 0, \quad (4.3)$$

$$\mathcal{J}_2^0 = 0. \quad (4.4)$$

Replacing equations (4.3-4.4) into eq. (2.14) yields a zero momentum, while by substituting eq.(4.2) into eq.(2.15), we get the energy of a massless scalar field contained in a “sphere” of radius $\mathcal{X}$ for the three-dimensional spacetime described by the line element(3.6)

$$E(\mathcal{X}) = \frac{4\pi}{K} \sqrt{c^2 + 4\lambda^2} (\cosh(2\mathcal{X}) - 1). \quad (4.5)$$

This is also the energy of the gravitational field that the scalar field experiences at a finite distance $\mathcal{X}$. Thus, the energy given by equation (4.5) is also called effective gravitational mass ($E = M_{\text{eff}}$) of the spacetime under study. The energy content of an infinitely large “sphere” is infinite, as it is also evident from Figure 1.




Figure 1: Energy versus distance

5 Discussion - Conclusions

As we pointed out previously, the subject of the localization of energy continues to be open, since Einstein, in his theory of special relativity, declared that mass is equivalent to energy. Misner *et al.* [20] concluded that the energy is localizable only for spherical systems. On the other hand, Cooperstock and Sarracino [21] demonstrated that if the energy is localizable in spherical systems then it is also localizable in any spacetime.

Now, concerning the use of several energy-momentum complexes for the localization of energy and momentum, we remark that although it has many adepts, there has also been a great deal of criticism related to the use of such prescriptions. The main disadvantage of the energy-momentum complexes lies in that most of them restrict one to make calculations in quasi-Cartesian coordinates. Nevertheless, Møller's energy-momentum complex is the only one allowing calculations in any coordinate system.

In this work, we have explicitly calculated the energy and momentum density distributions associated with an exact solution of Einstein's field equations with a negative cosmological constant and minimally coupled to a static massless scalar field in a static, spherically symmetric, three-dimensional spacetime. The energy and momentum distributions are computed by use of Møller's energy-momentum complex.

A physically acceptable expression for the energy density distribution is obtained. The momentum density distributions for the used $(2+1)$ -dimensional, static spherically symmetric background vanish. Additionally, the effective gravitational mass, i.e. the total energy, is explicitly evaluated for the specific gravitational background. We have shown that the energy distribution can be interpreted as the effective gravitational mass corresponding to the $(2+1)$ -dimensional background. The parameter $\mu \in (0, 1)$ appearing in (4.5) expresses the strength of the coupled scalar field. One can see that for the special case $\mu = 1$, the three-dimensional Anti-deSitter spacetime is obtained.

Furthermore, Møller's energy-momentum complex applied to this $(2+1)$ -dimensional solution satisfies the local conservation equation (2.13).

We point out that Møller's formalism provides physically satisfactory results for the aforementioned background. Finally, these results support Lessner's viewpoint [15] about the significance of Møller's energy-momentum complex as a powerful tool for the evaluation of the energy and momentum density distributions in a given geometry.

Acknowledgements

The authors are indebted to Dr. E.C. Vagenas for useful suggestions and comments.

References

- [1] A. Einstein, *Preuss. Akad. Wiss. Berlin* **47**, 778 (1915); Addendum-*ibid.* **47**, 799 (1915).
- [2] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields* (Pergamon Press, 1987, p. 280).
- [3] A. Papapetrou, *Proc. R. Irish. Acad.* **A52**, 11 (1948).
- [4] P. G. Bergmann and R. Thompson, *Phys. Rev.* **89**, 400 (1953).
- [5] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of General Theory of Relativity* (John Wiley and Sons, Inc., New York, 1972, p. 165).
- [6] C. Møller, *Ann. Phys. (NY)* **4**, 347 (1958).
- [7] K. S. Virbhadra, *Phys. Rev.* **D60**, 104041 (1999).
- [8] K. S. Virbhadra, *Phys. Rev.* **D41**, 1086 (1990); S. Virbhadra, *Phys. Rev.* **D42**, 2919 (1990); N. Rosen and K. S. Virbhadra, *Gen. Rel. Grav.* **25**, 429 (1993); K. S. Virbhadra and J. C. Parikh, *Phys. Lett.* **B331**, 302 (1994); K. S. Virbhadra, *Pramana - J. Phys.* **45**, 215 (1995); J. M. Aguirregabiria, A. Chamorro and K. S. Virbhadra, *Gen. Rel. Grav.* **28**, 1393 (1996).
- [9] A. Chamorro and K. S. Virbhadra, *Pramana - J. Phys.* **45**, 181 (1995); A. Chamorro and K. S. Virbhadra, *Int. J. Mod. Phys.* **D5**, 251 (1997); S. S. Xulu, *Int. J. Theor. Phys.* **37**, 1773 (1998); S. S. Xulu, *Int. J. Mod. Phys.* **D7**, 773 (1998); E. C. Vagenas, *Int.J.Mod.Phys.* **A18**, 5949 (2003) ; I. Radinschi, *Acta Physica Slovaca*, **49(5)**, 789 (1999); I. Radinschi, *Mod. Phys. Lett.* **A15**, Nos. 11&12, 803 (2000); I-Ching Yang and I. Radinschi, *Chin.J.Phys.* **41**, 326 (2003); I. Radinschi, *New Developments in String Theory Research*, Nova Science, 2005; T. Bringley, *Mod. Phys. Lett.* **A17**, 157 (2002); M. Súkeník and J. Sima, gr-qc/0101026; M. Sharif and Tasnim Fatima, gr-qc/0410004, accepted for publication in *Int. J. Mod. Phys. A*; M. Sharif, *Nuovo Cim.* **B19**, 463 (2004); M. Sharif, *Int.J.Mod.Phys.* **D13**, 1019 (2004); R. Gad, *Mod.Phys.Lett.* **A19**, 1847 (2004); R. Gad, *Astrophys.Space Sci.* **295**, 451 (2005); R. Gad, *Astrophys.Space Sci.* **293**, 453 (2004); O. Patashnick, gr-qc/0408086.
- [10] E. C. Vagenas, *Int.J.Mod.Phys.* **A18**, 5949 (2003); E. C. Vagenas, *Int.J.Mod.Phys.* **A18**, 5781 (2003); E. C. Vagenas, *Int.J.Mod.Phys.* **D14**, 573 (2005); Th. Grammenos, *Mod. Phys. Lett.*, **A20**, 1741 (2005).
- [11] F. I. Cooperstock, *Mod. Phys. Lett.* **A14**, 1531 (1999).
- [12] Chia-Chen Chang, J. M. Nester and Chiang-Mei Chen, *Phys. Rev. Lett.* **83**, 1897 (1999).

- [13] S. S. Xulu, *Mod. Phys. Lett.* **A15**, 1511 (2000); *PhD Thesis*, hep-th/0308070; S.S. Xulu, *Astrophys. Space. Sci.* **23**, (2003).; I-Ching Yang, Ching-Tzung Yeh, Rue-Ron Hsu and Chin-Rong Lee, *Int. J. Mod. Phys.* **D6**, 349 (1997); I-Ching Yang, Wei-Fui Lin and Rue-Ron Hsu, *Chin. J. Phys.* **37**, 113 (1999); E. C. Vagenas, *Int.J.Mod.Phys.* **A18**, 5781 (2003); E. C. Vagenas, *Mod.Phys.Lett.* **A19**, 213 (2004); R. Gad, gr-qc/0403039; R. Gad, *Astrophys.Space Sci.* **295**, 459 (2005); R. Gad, *Astrophys.Space Sci.* **293**, 453 (2004); I. Radinschi, *Mod. Phys. Lett.* **A15 (35)**, 2171 (2000); M. Sharif and Tasnim Fatima, gr-qc/0410004, accepted for publication in *Int. J. Mod. Phys. A*.
- [14] I. Radinschi, *Fizika B* **B9(4)**, 203 (2000); I. Radinschi, *Mod.Phys.Lett.* **A15**, 2171 (2000); I. Radinschi, *Chin. J. Phys.* **39(3)**, 231 (2001); I. Radinschi, *Mod. Phys. Lett.* **A16(10)**, 673 (2001); I-Ching Yang and I. Radinschi, *Chin.J.Phys.* **42**, 40 (2004); I-Ching Yang and I. Radinschi, *Chin.J.Phys.* **41**, 326 (2003); I-Ching Yang and I. Radinschi, *Mod. Phys. Lett.* **A17**, 1159 (2002); I. Radinschi and I-Ching Yang, gr-qc/0405118.
- [15] G. Lessner, *Gen. Relativ. Gravit.* **28**, 527 (1996).
- [16] H. Bondi, *Proc. R. Soc. London* **A427**, 249 (1990).
- [17] D. Daghan and A. H. Bilge, gr-qc/0505130.
- [18] H.-J. Matschull, *Class.Quant. Grav.* **16**, 1069 (1999).
- [19] R.M. Wald, *General Relativity* (University of Chicago Press, 1984, p.70).
- [20] C.W. Misner, K.S. Thorne and J.A. Wheeler, *Gravitation* (W.H. Freeman and Co., NY, 1973, p. 603).
- [21] F.I. Cooperstock and R.S. Sarracino, *J. Phys. A: Math. Gen.* **11**, 877 (1978).